

Scalars 2015

Contribution ID : 3

Asymptotically free scaling solutions in nonabelian Higgs models

Content :

We construct asymptotically free renormalization group trajectories for the generic nonabelian Higgs model in four-dimensional spacetime. These ultraviolet-complete trajectories become visible by generalizing the renormalization/boundary conditions in the definition of the correlation functions of the theory. Though they are accessible in a controlled weak-coupling analysis, these trajectories originate from threshold phenomena which are missed in a conventional perturbative analysis relying on the deep Euclidean region. We identify a candidate three-parameter family of renormalization group trajectories interconnecting the asymptotically free ultraviolet regime with a Higgs phase in the low-energy limit. We provide estimates of their low-energy properties in the light of a possible application to the standard model Higgs sector.

Primary authors : Prof. GIES, Holger (Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, D-07743 Jena, Germany) ; Dr. ZAMBELLI, Luca (Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, D-07743 Jena, Germany)

Co-authors :

Presenter : Dr. ZAMBELLI, Luca (Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, D-07743 Jena, Germany)

Session classification : --not yet classified--

Track classification : --not yet classified--

Type : --not specified--