

# Scalars 2015

Contribution ID : 49

## SIMP model at NNLO in chiral perturbation theory

### Content :

We investigate the phenomenological viability of a recently proposed class of composite dark matter models where the relic density is determined by  $3 \rightarrow 2$  number-changing processes in the dark sector. Here the pions of the strongly interacting field theory constitute the dark matter particles. By performing a consistent next-to-leading- and next-to-next-to-leading-order chiral perturbative investigation we demonstrate that the leading-order analysis cannot be used to draw conclusions about the viability of the model. We further show that higher-order corrections substantially increase the tension with phenomenological constraints challenging the viability of the simplest realization of the strongly interacting massive particle paradigm.

### Summary :

Phys.Rev. D92 (2015) 075036  
DOI: 10.1103/PhysRevD.92.075036  
arXiv:1507.01590

**Primary authors** : Mr. LANGÆBLE, Kasper (CP3-Origins) ; Prof. SANNINO, Francesco (CP3-origins, University of Southern Denmark) ; Mr. HANSEN, Martin (CP3-Origins)

**Co-authors** :

**Presenter** : Mr. LANGÆBLE, Kasper (CP3-Origins)

**Session classification** : --not yet classified--

**Track classification** : --not yet classified--

**Type** : --not specified--