

# Natural SUSY, Muon $g-2$ , and Higgs Measurements at Future $e^-e^+$ Colliders

Kechen Wang

DESY, Hamburg, Germany

20<sup>th</sup> Planck Conference, U of Warsaw  
May 25, 2017

Based on [Phys. Rev. D **93** (2016) 055040], with T. Li, S. Raza

# Outline

- ★ Parameter Space Scan
  - ◆ Introduction
  - ◆ Theoretical Dependences
  - ◆ Input Constraints
  
- ★ Numerical Results
  - ◆ Electroweak Fine-tuning
  - ◆ Higgs Coupling Measurements
  - ◆ Muon Magnetic Moment Measurement
  - ◆ Collider Phenomenology
  
- ★ Summary




# Current Status of BSM Searches

SUSY from LHC:

- $m_{\text{gluino}} > 1.9 \text{ TeV}$
- $m_{\text{stop}} > 950 \text{ GeV}$
- $m_{\text{chargino}} > 600 \text{ GeV}$

DM from direct detection [1608.07648]


→ lots of parameter space excluded



→ No clear Hints of New Physics so far.



# Current Status of BSM Searches



Perhaps we will have to face the “nightmare scenario” after LHC

→ no clear hints of BSM physics.

→ no definite guidance to BSM physics.

=> have to search for all possible directions.




# Two Approaches to Constrain BSM

Feasible step after LHC:

Higgs property (couplings, decay width, spin & CP) measurements @ e<sup>-</sup>e<sup>+</sup> colliders

=> new physics indirectly

## EFT



D>4 EFT Operators


(see talks of C. Grojean, T. Vantalón, J. Gu, ...)

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{SM}} + \sum_i \frac{1}{\Lambda^{d_i-4}} c_i \mathcal{O}_i$$

- ✓ Simple, Generic, model-independent
- ✗ Validation, may not be valid

## UV-complete Models

(see talks of M. Drees, A. Choudhury, ...)



SUSY, Composite, Extra-D, ...

- ✓ Concrete, more details & predicable
- ✗ Specified, model-dependent

This talk: scanning the parameter space of a natural SUSY model

← Higgs coupling & Muon anomalous magnetic moment measurements

# Theoretical Dependences

## Higgs Couplings in MSSM model

In the decoupling limit of  $m_A \gg m_Z$

$$k_V \equiv \frac{g_{hVV}^{\text{SUSY}}}{g_{hVV}^{\text{SM}}} = \sin(\alpha + \beta) \sim 1 - \mathcal{O}\left(\frac{m_Z^4}{m_A^4} \frac{2}{\tan^2 \beta}\right), \text{ for } V = W, Z$$

$$k_{b,\tau} \sim 1 - \mathcal{O}\left(\frac{m_Z^4}{m_A^4} \frac{2}{\tan^2 \beta}\right) + \mathcal{O}\left(\frac{m_Z^2}{m_A^2} \frac{2}{\tan \beta} \tan \beta\right) \sim 1 + \mathcal{O}\left(2 \frac{m_Z^2}{m_A^2}\right)$$

$$k_t \sim 1 - \mathcal{O}\left(\frac{m_Z^4}{m_A^4} \frac{2}{\tan^2 \beta}\right) - \mathcal{O}\left(\frac{m_Z^2}{m_A^2} \frac{2}{\tan \beta} \cot \beta\right) \sim 1 - \mathcal{O}\left(\frac{m_Z^2}{m_A^2} \frac{2}{\tan^2 \beta}\right)$$

$$k_g \approx 1 + \frac{m_t^2}{4} \left[ \frac{1}{m_{\tilde{t}_1}^2} + \frac{1}{m_{\tilde{t}_2}^2} - \frac{X_t^2}{m_{\tilde{t}_1}^2 m_{\tilde{t}_2}^2} \right] \quad X_t = |A_t - \mu / \tan \beta|$$

Muon anomalous magnetic moment from  $\tilde{e}_{R,L} - \tilde{\chi}_1^0$  and  $\tilde{\nu} - \tilde{\chi}_1^\pm$  loops

$$\Delta a_\mu^{\text{SUSY}} \sim \frac{M_i \mu \tan \beta}{m_{\text{SUSY}}^4}$$

$M_{1,2} \rightarrow$  weak-scale gaugino masses;  
 $\mu \rightarrow$  Higgsino mass parameter;  
 $m_{\text{SUSY}} \rightarrow$  sparticle mass circulating in the loop

# Input Constraints

Scan the parameter space of GmSUGRA model with  
small fine-tuning  $\Delta_{EW} \leq 100$

Using the **ISAJET** 7.85 package

from LEP2  $m_{\tilde{t}_1}, m_{\tilde{b}_1} \geq 100 \text{ GeV},$   
 $m_{\tilde{\tau}_1} \geq 105 \text{ GeV},$   
 $m_{\tilde{\chi}_1^\pm} \geq 103 \text{ GeV}.$

from LHC  $1.7 \text{ TeV} \leq m_{\tilde{g}} \text{ (for } m_{\tilde{g}} \sim m_{\tilde{q}})$   
 $1.3 \text{ TeV} \leq m_{\tilde{g}} \text{ (for } m_{\tilde{g}} \ll m_{\tilde{q}})$   
 $300 \text{ GeV} \leq m_A$

$$123 \text{ GeV} \leq m_h \leq 127 \text{ GeV}$$

$$1.6 \times 10^{-9} \leq \text{BR}(B_s \rightarrow \mu^+ \mu^-) \leq 4.2 \times 10^{-9} \text{ (} 2\sigma)$$

$$2.99 \times 10^{-4} \leq \text{BR}(b \rightarrow s\gamma) \leq 3.87 \times 10^{-4} \text{ (} 2\sigma)$$

$$0.70 \times 10^{-4} \leq \text{BR}(B_u \rightarrow \tau\nu_\tau) \leq 1.5 \times 10^{-4} \text{ (} 2\sigma)$$

## (I) Basic constraints:

neutralino LSP;

fine-tuning;

sparticle mass

Higgs mass;

B-physics.



# Input Constraints

## (II) Muon magnetic moment constraint

$$4.7 \times 10^{-10} \leq \Delta a_\mu \leq 52.7 \times 10^{-10} \quad (3\sigma)$$

## (III) Higgs couplings constraint

Collider	HL-LHC	ILC	CEPC (2 IP)	FCC-ee (4 IP)
$\sqrt{s}$ (GeV)	14000	500	240	240
$\mathcal{L}$ (fb <sup>-1</sup> )	3000	500	5000	10000
polarization ( $e^-, e^+$ )	-	(-0.8, +0.3)	(0, 0)	(0, 0)
$k_g$	9.1	2.3	1.5	1.1
$k_W$	5.1	1.2	1.2	0.85
$k_Z$	4.4	1.0	0.26	0.16
$k_\gamma$	4.9	8.4	4.7	1.7
$k_b$	12	1.7	1.3	0.88
$k_\tau$	9.7	2.4	1.4	0.94
$k_t$	11	14	-	-

Table 1: Summary for the precisions of Higgs boson coupling measurements in percentage at different colliders.

5 ~ 10 %


~1 %

For ILC, 250 GeV running & luminosity upgrade → better precisions.

# Outline





- ★ Parameter Space Scan
  - ◆ Introduction
  - ◆ Theoretical Dependences
  - ◆ Input Constraints
  
- ★ Numerical Results
  - ◆ Electroweak Fine-tuning
  - ◆ Higgs Coupling Measurements
  - ◆ Muon Magnetic Moment Measurement
  - ◆ Collider Phenomenology
  
- ★ Summary

# Electroweak Fine-tuning $\Delta_{EW}$






# Electroweak Fine-tuning $\Delta_{EW}$







Higgs mass constraint

Higgs coupling constraint

# Higgs Coupling Measurements




# Higgs Coupling Measurements






# Muon Magnetic Moment Measurement




# Muon Magnetic Moment Measurement

All points – (I) + (II) + (III) Higgs couplings @ CEPC



# Collider Phenomenology




Precisions of  $Hbb$  coupling: > 10%, 5%~10%, 3%~5%, < 3%





→ Complementary to the direct collider searches



# Collider Phenomenology



# Collider Phenomenology



# Benchmark Points

	large Higgs deviations	large $\Delta a_\mu$	A resonance	stau-neutralino coannihilation
	Point 1	Point 2	Point 3	Point 4
$m_{0U}$	1037	2582	5057	5327
$m_{\tilde{Q}}$	1003.3	2366.4	4631	4879.2
$m_{\tilde{U}^c}$	1162.1	3306.7	6487	6830.6
$m_{\tilde{D}^c}$	1311.7	3326.3	6474.9	6841.3
$m_{\tilde{L}}$	328.1	264.8	1023	858.2
$m_{\tilde{E}^c}$	814.1	514.8	900.9	978.6
$M_1$	857.1	350.9	602.7	686.5
$M_2$	656	993	634.3	828.8
$M_3$	1158.8	-612.25	555.3	473.05
$A_t = A_b$	-3292	5390	-9095	-9684
$A_{\tilde{\tau}}$	-691.6	1192	-1570	-2937
$\tan \beta$	11.6	33.8	19.7	21.4
$\mu$	183	172.5	377.5	168.2
$m_A$	<u>338.9</u>	754.9	<u>603</u>	846
$\Delta_{\text{HS}}$	1883	3151	10873	11993.67
$\Delta_{\text{EW}}$	27	23	87	87
$\Delta a_\mu$	$4.731 \times 10^{-10}$	<u><math>16.63 \times 10^{-10}</math></u>	$7.683 \times 10^{-10}$	$12.161 \times 10^{-10}$
$m_h$	123	123	125	125
$m_H$	342	751	607	856
$m_{H^\pm}$	348	750	608	850
$\kappa_b, \kappa_t$	<u>1.22541</u> , 0.99817	1.04295, 0.99996	1.06459, 0.99983	1.03296, 0.99992
$\kappa_\tau, \kappa_W = \kappa_Z$	<u>1.22756</u> , 0.99981	1.04787, 0.99999	1.06644, 0.99999	1.03454, 0.99999
$\kappa_g, \kappa_\gamma$	0.99083, 1.00369	1.00473, 0.99763	0.99954, 1.00008	1.00038, 0.99714
$m_{\tilde{\chi}_{1,2}^0}$	171, 191	131, 183	<u>260</u> , 376	<u>161</u> , 178
$m_{\tilde{\chi}_{3,4}^0}$	368, 541	195, 834	390, 567	312, 717
$m_{\tilde{\chi}_{1,2}^\pm}$	185, 534	179, 823	378, 555	175, 702
$m_{\tilde{g}}$	2548	1539	1498	1317
$m_{\tilde{u}_{L,R}}$	2419, 2558	2705, 3476	4721, 6513	4942, 6832
$m_{\tilde{t}_{1,2}}$	1036, 1798	1151, 1762	2311, 3401	2326, 3543
$m_{\tilde{d}_{L,R}}$	2421, 2532	2706, 3547	4722, 6595	4943, 6938
$m_{\tilde{b}_{1,2}}$	1771, 2468	1212, 3089	2380, 6340	2401, 6630
$m_{\tilde{\nu}_{1,2}}$	726	<u>521</u>	698	507
$m_{\tilde{\nu}_3}$	718	343	645	308
$m_{\tilde{e}_{L,R}}$	737, 525	<u>514</u> , 787	651, 1409	398, 1487
$m_{\tilde{\tau}_{1,2}}$	518, 728	353, 568	627, 1338	<u>185</u> , 1333
$\Omega_{\text{CDM}} h^2$	0.0014	0.0846	<u>0.1017</u>	0.0099

# Summary



- ★ Higgs & Other Low Energy Measurements => BSM physics
  - If no clear hints of BSM after LHC
  - EFT or a UV-complete model
  
- ★ Scanning Parameter Space of a Natural SUSY Model
  - Using Higgs coupling @ HL-LHC, ILC, CEPC & muon g-2
  - $\Delta_{EW} \sim 30$
  - $m_A > 1.2$  (1.1) TeV from  $k_b$  ( $k_T$ ) at CEPC
  - $\tilde{l}_R : [0.6 - 2]$  TeV,  $\tilde{l}_L, \tilde{\nu} : [0.4 - 1.2]$  TeV with  $\tilde{\chi}_1^0 : < 400$  GeV from  $\Delta a_\mu$
  - $\tan\beta$  vs.  $m_A$ : complementary to the direct collider searches
  - 4 Benchmark points


The background features several faint, overlapping postmarks and stamps in a light brown or tan color. These include circular postmarks with dates and times, and rectangular stamps with text like 'POSTAGE' and 'AIR MAIL'. The overall aesthetic is that of an old, weathered document or envelope.

*Thank you for your attention !*

*Any Questions ?*

# Backup Slide

## Higgs Coupling Measurements



$k_\gamma < 1$   
← negative contributions from sbottom & stau



# Backup Slide

## General Minimal Supergravity (GmSUGRA) Model

### Scanning the parameter space

$$0 \text{ GeV} \leq m_0^U \leq 9000 \text{ GeV},$$

$$100 \text{ GeV} \leq M_1 \leq 2000 \text{ GeV},$$

$$100 \text{ GeV} \leq M_2 \leq 2100 \text{ GeV},$$

$$100 \text{ GeV} \leq m_{\tilde{L}} \leq 1200 \text{ GeV},$$

$$100 \text{ GeV} \leq m_{\tilde{E}^c} \leq 1200 \text{ GeV},$$

$$100 \text{ GeV} \leq \mu \leq 1500 \text{ GeV},$$

$$0 \text{ GeV} \leq m_A \leq 9500 \text{ GeV},$$

$$-16000 \text{ GeV} \leq A_U = A_D \leq 18000 \text{ GeV}$$

$$-6000 \text{ GeV} \leq A_E \leq 6000 \text{ GeV},$$

$$2 \leq \tan\beta \leq 60.$$

→ sleptons, charginos, bino, wino,  
Higgsinos within one TeV;

→ squarks, gluinos can be in several TeV mass

# Backup Slide

## Electroweak Fine-tuning

Use ISAJET 7.85 to calculate the fine-tuning measure at the EW scale  $M_{EW}$   
→ lower values of  $\Delta_{EW}$  corresponds to less fine-tuning

Z boson mass  $M_Z$ , after including the one-loop effective potential contributions to the tree-level MSSM Higgs potential

$$\frac{m_Z^2}{2} = \frac{m_{H_d}^2 + \Sigma_d^d - (m_{H_u}^2 + \Sigma_u^u) \tan^2 \beta}{\tan^2 \beta - 1} - \mu^2$$

$$\Delta_{EW} \equiv \max(C_k) / (M_Z^2/2)$$