

Supergeometric Scalars and Fermions

APOSTOLOS PILAFTSIS

*Department of Physics and Astronomy, University of Manchester,
Manchester M13 9PL, United Kingdom*

SCALARS 2023

Warsaw, Poland, September 2023

Based on: K Finn, S Karamitsos, AP, PRD102 (2020) 045014 [arXiv:1910.06661]
EPJC81 (2021) 572 [arXiv:2006.05831]

$\Rightarrow$ Viola Gattus, AP, arXiv:2307.01126 and arXiv:2310.nnnnn

Related works: S Karamitsos, AP, NPB907 (2016) 785; NPB927 (2018) 219

Outline:

- Brief History on Covariant or Geometric Methods in QFT
- Grand Covariance in Quantum Gravity
- The Fermion problem and Supermanifolds
- Supergeometric Effective Actions for Scalars and Fermions
- Conclusions & Future Directions

• Brief History on Covariant Methods in QFT

- B DeWitt '67: . . . *The Manifestly Covariant Quantum Field Theory*, introducing the well-known *Background Field Method*.
- S Weinberg '68, . . . J Honerkamp '72 . . . :
Geometry of Field Space and Amplitudes in Non-linear σ -Models
 $\Rightarrow$ Little Higgs Models, SMEFT $\subset$ HEFT, Multifield Inflation . . .
- L Alvarez-Gaumé, DZ Friedman, S Mukhi '81:
Geometry and UV Finiteness of SUSY Non-linear σ -Models
- G Vilkovisky '84, B DeWitt '85,
Gauge Independence of Covariant VDW Effective Action
- MK Gaillard '86: *Gauge-Covariant Derivative Approach to EFT*
 $\Rightarrow$ Universal effective action, threshold effects in SMEFT . . .

- JM Cornwall '82, JM Cornwall, J Papavassiliou '89, AP '97:
Pinch Technique: a Gauge-Covariant Diagrammatic Approach to Off-Shell Amplitudes
⇒ Effective gluon mass, Non-Abelian effective charges,
Unstable particle dynamics and High-energy unitarity of Amplitudes
- G Sigl, G Raffelt '93, PSB Dev, P Millington, AP, D Teresi '14:
Flavour Covariant Transport Equations
⇒ Neutrino flavour dynamics in astrophysics and cosmology,
Leptogenesis . . .
- D Teresi, AP '13:
Symmetry Improved Cornwall-Jackiw-Tomboulis Effective Action
⇒ Massless Goldstones, 2nd order phase transitions for $\mathbb{O}(N)$ theories,
RG exactness and IR safe effective potentials . . .

⋮

• Grand Covariance in Quantum Gravity

[G Vilkovisky '84, B DeWitt, '85;
K Finn, S Karamitsos, AP, '19]

$$S = \int d^D x \sqrt{-g} \left[-\frac{f(\varphi)}{2} R + \frac{1}{2} k_{AB}(\varphi) g^{\mu\nu} (\nabla_\mu \varphi^A) (\nabla_\nu \varphi^B) - V(\varphi) \right],$$

$S = S[g_{\mu\nu}, \varphi; f(\varphi), k(\varphi), V(\varphi)]$: *classical action*
 $f(\varphi), k_{AB}(\varphi), V(\varphi)$: *model functions*

• Grand Covariance in Quantum Gravity

[G Vilkovisky '84, B DeWitt, '85;
K Finn, S Karamitsos, AP, '19]

$$S = \int d^D x \sqrt{-g} \left[-\frac{f(\varphi)}{2} R + \frac{1}{2} k_{AB}(\varphi) g^{\mu\nu} (\nabla_\mu \varphi^A) (\nabla_\nu \varphi^B) - V(\varphi) \right],$$

$$S = S[g_{\mu\nu}, \varphi; f(\varphi), k(\varphi), V(\varphi)]: \text{classical action}$$

$$f(\varphi), k_{AB}(\varphi), V(\varphi): \text{model functions}$$

Grand or Frame Covariance:

(i) *Spacetime diffeomorphisms*

$$x^\mu \rightarrow \tilde{x}^\mu = \tilde{x}^\mu(x^\nu), \quad \text{with} \quad ds^2 = g_{\mu\nu} dx^\mu dx^\nu = \tilde{g}_{\mu\nu} d\tilde{x}^\mu d\tilde{x}^\nu$$

(ii) *Field reparametrizations*

$$g_{\mu\nu} \rightarrow \tilde{g}_{\mu\nu} = \tilde{g}_{\mu\nu}(g_{\kappa\lambda}, \varphi) = \Omega^2(\varphi) g_{\mu\nu}$$

$$\varphi^A \rightarrow \tilde{\varphi}^A = \tilde{\varphi}^A(g_{\mu\nu}, \varphi) = \tilde{\varphi}^A(\varphi)$$

• Grand Covariance in Quantum Gravity

[G Vilkovisky '84, B DeWitt, '85;
K Finn, S Karamitsos, AP, '19]

$$S = \int d^D x \sqrt{-g} \left[-\frac{f(\varphi)}{2} R + \frac{1}{2} k_{AB}(\varphi) g^{\mu\nu} (\nabla_\mu \varphi^A) (\nabla_\nu \varphi^B) - V(\varphi) \right],$$

$$S = S[g_{\mu\nu}, \varphi; f(\varphi), k(\varphi), V(\varphi)]: \text{classical action}$$

$$f(\varphi), k_{AB}(\varphi), V(\varphi): \text{model functions}$$

Grand or Frame Covariance:

(i) *Spacetime diffeomorphisms*

$$x^\mu \rightarrow \tilde{x}^\mu = \tilde{x}^\mu(x^\nu), \quad \text{with} \quad \underbrace{ds^2 = g_{\mu\nu} dx^\mu dx^\nu \equiv \tilde{g}_{\mu\nu} d\tilde{x}^\mu d\tilde{x}^\nu}_{\text{Not invariant under (ii)}}$$

(ii) *Field reparametrizations*

$$g_{\mu\nu} \rightarrow \tilde{g}_{\mu\nu} = \tilde{g}_{\mu\nu}(g_{\kappa\lambda}, \varphi) = \underline{\Omega^2(\varphi) g_{\mu\nu}}$$

$$\varphi^A \rightarrow \tilde{\varphi}^A = \tilde{\varphi}^A(g_{\mu\nu}, \varphi) = \tilde{\varphi}^A(\varphi)$$

- Introduce **new** model function $\ell(\varphi)$ to restore (i)

$$ds^2 = \bar{g}_{\mu\nu} dx^\mu dx^\nu, \quad \text{with} \quad \bar{g}_{\mu\nu} \equiv \frac{g_{\mu\nu}}{\ell^2(\varphi)}$$

- Introduce **new** model function $\ell(\varphi)$ to restore (i)

$$ds^2 = \bar{g}_{\mu\nu} dx^\mu dx^\nu, \quad \text{with} \quad \bar{g}_{\mu\nu} \equiv \frac{g_{\mu\nu}}{\ell^2(\varphi)}$$

- Transformation of model functions

$$\tilde{\ell}(\varphi) = \Omega \ell(\varphi),$$

$$\tilde{f}(\varphi) = \Omega^{-2} f(\varphi),$$

$$\tilde{k}_{\tilde{A}\tilde{B}}(\varphi) = \left[k_{AB} - 6f(\ln \Omega)_{,A}(\ln \Omega)_{,B} + 3f_{,A}(\ln \Omega)_{,B} + 3(\ln \Omega)_{,A}f_{,B} \right] \partial^A \varphi_{\tilde{A}} \partial^B \varphi_{\tilde{B}},$$

$$\tilde{V}(\varphi) = \Omega^{-4} V(\varphi).$$

- Introduce **new** model function $\ell(\varphi)$ to restore (i)

$$ds^2 = \bar{g}_{\mu\nu} dx^\mu dx^\nu, \quad \text{with} \quad \bar{g}_{\mu\nu} \equiv \frac{g_{\mu\nu}}{\ell^2(\varphi)}$$

- Transformation of model functions

$$\tilde{\ell}(\varphi) = \Omega \ell(\varphi),$$

$$\tilde{f}(\varphi) = \Omega^{-2} f(\varphi),$$

$$\tilde{k}_{\tilde{A}\tilde{B}}(\varphi) = \left[k_{AB} - 6f(\ln \Omega)_{,A}(\ln \Omega)_{,B} + 3f_{,A}(\ln \Omega)_{,B} + 3(\ln \Omega)_{,A}f_{,B} \right] \partial^A \varphi_{\tilde{A}} \partial^B \varphi_{\tilde{B}},$$

$$\tilde{V}(\varphi) = \Omega^{-4} V(\varphi).$$

- Frame invariance of the classical action S :

$$S[\tilde{g}_{\mu\nu}, \tilde{\varphi}; \tilde{\ell}(\varphi), \tilde{f}(\varphi), \tilde{k}(\varphi), \tilde{V}(\varphi)] = S[g_{\mu\nu}, \varphi; \ell(\varphi), f(\varphi), k(\varphi), V(\varphi)]^*$$

Models related by a frame transformation define an *equivalence class*

[K Falls, M Herrero-Valea '19]

* $S = \int d^D x \sqrt{-g} \mathcal{L} = \int d^D x \sqrt{-\bar{g}} \bar{\mathcal{L}}$ is independent of $\ell(\varphi)$ (only at tree level).

– Coordinates of the **Grand** Configuration Space

$$\Phi^i \equiv \Phi^I(x_I) = \begin{pmatrix} g^{\mu\nu}(x) \\ \phi^A(x_A) \end{pmatrix}, \text{ with } i = \{I, x_I\}, I = \{\mu\nu, A\}, x_I = \{x, x_A\}.$$

– Coordinates of the **Grand** Configuration Space

$$\Phi^i \equiv \Phi^I(x_I) = \begin{pmatrix} g^{\mu\nu}(x) \\ \phi^A(x_A) \end{pmatrix}, \text{ with } i = \{I, x_I\}, I = \{\mu\nu, A\}, x_I = \{x, x_A\}.$$

– The **Grand** Configuration Space Metric

$$\mathcal{G}_{ij} \equiv \frac{\bar{g}_{\mu\nu}}{D} \frac{\bar{\delta}^2 S}{\bar{\delta}(\partial_\mu \Phi^i) \bar{\delta}(\partial_\nu \Phi^j)} = \ell^2 \begin{pmatrix} f P_{\mu\nu\rho\sigma} & -\frac{3}{4} f_{,B} g_{\mu\nu} \\ -\frac{3}{4} f_{,A} g_{\rho\sigma} & k_{AB} \end{pmatrix} \bar{\delta}^{(D)}(x_I - x_J),$$

where $\bar{\delta}^{(D)}(x_I - x_J) \equiv \delta^{(D)}(x_I - x_J) / \sqrt{-\bar{g}}$ is *frame invariant*, and

$$P_{\mu\nu\rho\sigma} \equiv G_{(\mu\nu)(\rho\sigma)} = \frac{1}{2} \left(g_{\mu\rho} g_{\sigma\nu} + g_{\mu\sigma} g_{\rho\nu} - \alpha g_{\mu\nu} g_{\rho\sigma} \right)$$

is the *gravitational field-space metric*.

Condition on the **inverse** metric $G^{(\mu\nu)(\rho\sigma)}$:

$$G^{(\mu\nu)(\rho\sigma)} = g^{\alpha\mu} g^{\beta\nu} g^{\kappa\rho} g^{\lambda\sigma} G_{(\alpha\beta)(\kappa\lambda)} \implies \alpha = 0 \text{ or } 1.$$

– Quantum Effective Action

$$\exp\left(\frac{i}{\hbar}\Gamma[\varphi]\right) = \int [\mathcal{D}\phi] \exp\left\{\frac{i}{\hbar}\left[S[\phi] + \frac{\delta\Gamma[\varphi]}{\delta\varphi^a}(\varphi^a - \phi^a)\right]\right\}.$$

– Quantum Effective Action

$$\exp\left(\frac{i}{\hbar}\Gamma[\varphi]\right) = \int [\mathcal{D}\phi] \exp\left\{\frac{i}{\hbar}\left[S[\phi] + \frac{\delta\Gamma[\varphi]}{\delta\varphi^a}(\varphi^a - \phi^a)\right]\right\}.$$

Not invariant under frame transformations

– Quantum Effective Action

$$\exp \left(\frac{i}{\hbar} \Gamma[\varphi] \right) = \int [\mathcal{D}\phi] \exp \left\{ \frac{i}{\hbar} \left[S[\phi] + \frac{\delta \Gamma[\varphi]}{\delta \varphi^a} (\varphi^a - \phi^a) \right] \right\}.$$

Not invariant under frame transformations.

– VDW Quantum Effective Action

$$\exp \left(\frac{i}{\hbar} \Gamma[\varphi] \right) = \int [\overline{\mathcal{D}}\Phi] \mathcal{M}[\Phi] \exp \left\{ \frac{i}{\hbar} \left[S[\Phi] + \frac{\delta \Gamma[\varphi]}{\delta \varphi^i} \Sigma^i[\varphi, \Phi] \right] \right\},$$

with $\varphi = (g^{\mu\nu}, \phi)$,

$$[\overline{\mathcal{D}}\Phi] = \exp \left[\sum_I \int d^D x \sqrt{-\bar{g}(x)} \ln \mathcal{D}\Phi^I(x) \right], \quad \mathcal{M}[\Phi] = V_{\text{FP}} \sqrt{\det(\mathcal{G}_{ij})},$$

and V_{FP} is the *Faddeev–Popov determinant* [for $\text{SU}(N)$, see Rebhan '87].

– One- and Two-Loop **VDW** Effective Actions

[consistent with P Ellicott, T Toms '89]

$$\Gamma^{(1)}[\varphi] = \frac{i}{2} \ln \overline{\det}(\nabla^a \nabla_b S),$$

$$\begin{aligned} \Gamma^{(2)}[\varphi] &= \text{Diagram 1} + \text{Diagram 2} \\ &= -\frac{1}{8} \Delta^{ab} \Delta^{cd} \nabla_{(a} \nabla_b \nabla_c \nabla_{d)} S \\ &\quad + \frac{1}{12} \Delta^{ab} \Delta^{cd} \Delta^{ef} (\nabla_{(a} \nabla_c \nabla_{e)} S) (\nabla_{(b} \nabla_d \nabla_{f)} S), \end{aligned}$$

with $\Delta^{ab} \equiv (\nabla_a \nabla_b S)^{-1} = \Delta^{ba}$.

– One- and Two-Loop **VDW** Effective Actions

[consistent with P Ellicott, T Toms '89]

$$\Gamma^{(1)}[\varphi] = \frac{i}{2} \ln \overline{\det}(\nabla^a \nabla_b S),$$

$$\begin{aligned} \Gamma^{(2)}[\varphi] &= \text{Diagram 1} + \text{Diagram 2} \\ &= -\frac{1}{8} \Delta^{ab} \Delta^{cd} \nabla_{(a} \nabla_b \nabla_c \nabla_{d)} S \\ &\quad + \frac{1}{12} \Delta^{ab} \Delta^{cd} \Delta^{ef} (\nabla_{(a} \nabla_c \nabla_{e)} S) (\nabla_{(b} \nabla_d \nabla_{f)} S), \end{aligned}$$

with $\Delta^{ab} \equiv (\nabla_a \nabla_b S)^{-1} = \Delta^{ba}$.

– **Grand** Covariance of the **VDW** Effective Action

[K Finn, S Karamitsos, AP, '19]

$$\Gamma[\varphi; \ell(\phi), f(\phi), k_{AB}(\phi), V(\phi)] = \Gamma[\tilde{\varphi}(\varphi); \tilde{\ell}(\phi), \tilde{f}(\phi), \tilde{k}_{AB}(\phi), \tilde{V}(\phi)]$$

with $\varphi = (g^{\mu\nu}, \phi)$.

– Recent works

- J. Honerkamp, Nucl. Phys. B36 (1972) 130; G. Ecker, J. Honerkamp, Phys. Lett. B42 (1972) 253.
- $\vdots$
- A. Y. Kamenshchik, C. F. Steinwachs, *Question of quantum equivalence between Jordan frame and Einstein frame*, Phys. Rev. D91 (2015) 084033.
- D. Burns, S. Karamitsos, AP, *Frame-Covariant Formulation of Inflation in Scalar-Curvature Theories*, Nucl. Phys. B907 (2016) 785.
- R. Alonso, E. Jenkins, A. Manohar, *Geometry in the Scalar Sector*, JHEP 08 (2016) 101
- S. Karamitsos, AP, *Frame Covariant Nonminimal Multifield Inflation*, Nucl. Phys. B927 (2018) 219.
- K. Finn, S. Karamitsos, AP, *Frame Covariance in Quantum Gravity*, Phys. Rev. D102 (2020) 045014
- R. Nagai, M. Tanabashi, K. Tsumura, Y. Uchida, Phys. Rev. D100 (2019) 075020
- A. Helset, A. Martin, M. Trott, JHEP03 (2020) 163
- K. Finn, S. Karamitsos, AP, *Frame covariant formalism for fermionic theories*, Eur. Phys. J. C81 (2021) 572
- T. Cohen, N. Craig, X. Lu, D. Sutherland, JHEP12 (2021) 003
- J. Talbert, JHEP 01 (2023) 069
- A. Helset, E.E. Jenkins, A.V. Manohar, *Geometry in scattering amplitudes*, Phys. Rev. D106 (2022) 116018
- V. Gattus and AP, *Minimal Supergeometric Quantum Field Theories*, arXiv:2307.01126.

- The **Fermion** problem and **Supermanifolds**


The Effective Action

Bryce De Witt '85

14 DISCUSSION

This completes the basic outline of how Vilkovisky's idea for an invariant scalar effective action works, or can be made to work. In discussing the advantages of such an effective action, I shall take up some of its possible defects. First of all, how unique is it? I do not think it is *in principle* as unique as Vilkovisky has claimed. Choices are made for three quantities: the starting metric γ_{IJ} , the functional measure $\mu_K[I, K]$; all else follows from these. The last two have no effect on the final form of Γ . The measure μ_I , and hence μ , is determined by unitarity requirements. Expression (13.11) for μ appears to depend on a fourth arbitrary quantity, g^+ , but in fact is independent of g^+ . To show this just vary ${}_af_\beta$ and use (13.15).

That leaves γ_{IJ} . Vilkovisky (1984) has suggested that γ_{IJ} should be determined by the coefficient of the highest derivatives in the superclassical field equations. This cannot be correct in the fermion sector of supergravity theory, where the highest-order derivative is first order, because the coefficient of a first-order derivative cannot yield a tensor of even rank having



- The **Fermion** problem and **Supermanifolds**


The Effective Action

Bryce De Witt '85

14 DISCUSSION

This completes the basic outline of how Vilkovisky's idea for an invariant scalar effective action works, or can be made to work. In discussing the advantages of such an effective action, I shall take up some of its possible defects. First of all, how unique is it? I do not think it is *in principle* as unique as Vilkovisky has claimed. Choices have been made for three quantities: the starting metric γ_{IJ} , the functional measure $\mu_K[I, K]$; all else follows from these. The last two have no effect on the final form of Γ . The measure μ_I , and hence μ , is determined by unitarity requirements. Expression (13.11) for μ appears to depend on a fourth arbitrary quantity, g^+ , but in fact is independent of g^+ . To show this just vary ${}_af_\beta$ and use (13.15).

That leaves γ_{IJ} . Vilkovisky (1984) has suggested that γ_{IJ} should be determined by the coefficient of the highest derivatives in the superclassical field equations. This cannot be correct in the fermion sector of supergravity theory, where the highest-order derivative is first order, because the coefficient of a first-order derivative cannot yield a tensor of even rank having



$\Rightarrow$ **No metric for non-SUSY theories with **fermions**:** $g_{XY}(\phi) \bar{\psi}^X i\gamma^\mu \partial_\mu \psi^Y$

- **Supergeometry on a Supermanifold** [K Finn, S Karamitsos, AP, EPJC81 (2021) 572;
V Gattus, AP, arXiv:2307.01126]

– **Fermions as Coordinates in the Field-Space Supermanifold**

$$\Phi \equiv \{\Phi^\alpha\} = (\phi^A, \psi_a^1, \bar{\psi}_{\dot{a}}^1, \psi_a^2, \bar{\psi}_{\dot{a}}^2, \dots)^T,$$

where ϕ^A are scalars and ψ_a^X are Dirac (or Weyl) fermions.

- **Supergeometry on a Supermanifold** [K Finn, S Karamitsos, AP, EPJC81 (2021) 572;
V Gattus, AP, arXiv:2307.01126]

– **Fermions as Coordinates in the Field-Space Supermanifold**

$$\Phi \equiv \{\Phi^\alpha\} = (\phi^A, \psi_a^1, \bar{\psi}_{\dot{a}}^1, \psi_a^2, \bar{\psi}_{\dot{a}}^2, \dots)^T,$$

where ϕ^A are scalars and ψ_a^X are Dirac (or Weyl) fermions.

– **Frame-covariant Lagrangian of a scalar theory with fermions**

$$\mathcal{L} = \underbrace{\frac{1}{2} g^{\mu\nu} \partial_\mu \Phi^\alpha \, {}_\alpha k_\beta(\Phi) \, \partial_\nu \Phi^\beta}_{: \text{scalars}} + \underbrace{\frac{i}{2} \zeta_\alpha^\mu(\Phi) \, \partial_\mu \Phi^\alpha}_{: \text{fermions}} - U(\Phi).$$

Model functions:

${}_\alpha k_\beta(\Phi)$: *rank-2 tensor* of the would-be bosonic metric (with ${}_\alpha k_X = 0$)

$\zeta_\alpha^\mu(\Phi)$: mixed spacetime and field-space *vector*

$U(\Phi)$: a *scalar* describing the potential and Yukawa sector

– Extracting the model functions ${}_{\alpha}k_{\beta}$ and ζ_{α}^{μ}

$${}_{\alpha}k_{\beta} = \frac{g_{\mu\nu}}{D} \frac{\overrightarrow{\partial}}{\partial(\partial_{\mu}\Phi^{\alpha})} \mathcal{L} \frac{\overleftarrow{\partial}}{\partial(\partial_{\nu}\Phi^{\beta})}, \quad \zeta_{\alpha}^{\mu} = \frac{2}{i} \left(\mathcal{L} - \frac{1}{2} g^{\mu\nu} \partial_{\mu}\Phi^{\alpha} {}_{\alpha}k_{\beta} \partial_{\nu}\Phi^{\beta} \right) \frac{\overleftarrow{\partial}}{\partial(\partial_{\mu}\Phi^{\alpha})}$$

– Extracting the model functions ${}_{\alpha}k_{\beta}$ and ζ_{α}^{μ}

$${}_{\alpha}k_{\beta} = \frac{g_{\mu\nu}}{D} \frac{\overrightarrow{\partial}}{\partial(\partial_{\mu}\Phi^{\alpha})} \mathcal{L} \frac{\overleftarrow{\partial}}{\partial(\partial_{\nu}\Phi^{\beta})}, \quad \zeta_{\alpha}^{\mu} = \frac{2}{i} \left(\mathcal{L} - \frac{1}{2} g^{\mu\nu} \partial_{\mu} \Phi^{\alpha} {}_{\alpha}k_{\beta} \partial_{\nu} \Phi^{\beta} \right) \frac{\overleftarrow{\partial}}{\partial(\partial_{\mu}\Phi^{\alpha})}$$

– Free theory as an example

$$\begin{aligned} \mathcal{L} = & \sum_{A \in N_{\text{scalars}}} \frac{1}{2} g^{\mu\nu} \partial_{\mu} \phi^A \partial_{\nu} \phi^A - \frac{1}{2} m_A^2 (\phi^A)^2 \\ & + \sum_{X \in M_{\text{fermions}}} \frac{i}{2} \left(\bar{\psi}^X \gamma^{\mu} \partial_{\mu} \psi^X - \partial_{\mu} \bar{\psi}^X \gamma^{\mu} \psi^X \right) - m_X \bar{\psi}^X \psi^X. \end{aligned}$$

Model functions:

$$\begin{aligned} {}_{\alpha}k_{\beta} &= \begin{pmatrix} \delta_{AB} & \mathbf{0}_{N \times 8M} \\ \mathbf{0}_{8M \times N} & \mathbf{0}_{8M \times 8M} \end{pmatrix}, \\ \zeta_{\alpha}^{\mu} &= \left(\mathbf{0}_N, \bar{\psi}_{\dot{a}}^1 \gamma_{\dot{a}a}^{\mu}, \gamma_{\dot{a}a}^{\mu} \psi_a^1, \bar{\psi}_{\dot{b}}^2 \gamma_{\dot{b}b}^{\mu}, \gamma_{\dot{b}b}^{\mu} \psi_b^2, \dots \right) \end{aligned}$$

– Deriving the Field-Space Metric

Define the rank-1 field-superspace tensor,

[improved by V Gattus, AP, arXiv:2307.01126]

$$\zeta_\alpha(\Phi) = \frac{1}{4} \frac{\delta \zeta_\alpha^\mu(\Phi)}{\delta \gamma^\mu} \xrightarrow{\text{improved}} \zeta_\alpha(\Phi) = \zeta_\beta^\mu(\Phi) \overleftarrow{\Sigma}_{\mu,\alpha}^\beta,$$

to derive the rank-2 anti-supersymmetric tensor (in analogy to $F_{\mu\nu}$ in QED)

$${}_\alpha \lambda_\beta(\Phi) = \frac{1}{2} \left(\frac{\overrightarrow{\partial}}{\partial \Phi^\alpha} \zeta_\beta(\Phi) - (-1)^{\alpha+\beta+\alpha\beta} \frac{\overrightarrow{\partial}}{\partial \Phi^\beta} \zeta_\alpha(\Phi) \right), \quad \text{with } \lambda^{\text{ST}} = -\lambda.$$

Introduce the *non*-singular rank-2 tensor:

$${}_\alpha \Lambda_\beta \equiv {}_\alpha k_\beta + {}_\alpha \lambda_\beta \xrightarrow{\text{free theory}} {}_\alpha N_\beta \equiv \begin{pmatrix} 1_N & 0 & 0 & 0 & 0 & \cdots \\ 0 & 0 & 1_4 & 0 & 0 & \cdots \\ 0 & 1_4 & 0 & 0 & 0 & \cdots \\ 0 & 0 & 0 & 0 & 1_4 & \cdots \\ 0 & 0 & 0 & 1_4 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

– Properties of the Field-Space **Super-Metric** ${}_{\alpha}G_{\beta}(\Phi)$

The **Super-Metric** ${}_{\alpha}G_{\beta}(\Phi)$ should:

1. Be *uniquely* determined from the *action*.
2. Transform as a proper *rank-2 field-space tensor*.
3. Be *supersymmetric* and *non-singular* to produce a non-zero line element.
4. Be *ultralocal*, i.e. it should not depend on $\partial_{\mu}\Phi$.
5. Have the *local form* on each point of the field-space Supermanifold

$${}_a H_b \equiv \begin{pmatrix} 1_N & 0 & 0 & 0 & 0 & \cdots \\ 0 & 0 & 1_4 & 0 & 0 & \cdots \\ 0 & -1_4 & 0 & 0 & 0 & \cdots \\ 0 & 0 & 0 & 0 & 1_4 & \cdots \\ 0 & 0 & 0 & -1_4 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

– The Field-Space **Super-Metric**

Determine first the *field-space vielbeins* ${}_{\alpha}e^a(\Phi)$ from

$${}_{\alpha}\Lambda_{\beta}(\Phi) = {}_{\alpha}e^a(\Phi) {}_aN_b {}^be_{\beta}^{s\top}(\Phi),$$

and use these to obtain the **Field-Space Super-Metric**:

$${}_{\alpha}G_{\beta}(\Phi) = {}_{\alpha}e^a(\Phi) {}_aH_b {}^be_{\beta}^{s\top}(\Phi)$$

– The Field-Space **Super-Metric**

Determine first the *field-space vielbeins* ${}_{\alpha}e^a(\Phi)$ from

$${}_{\alpha}\Lambda_{\beta}(\Phi) = {}_{\alpha}e^a(\Phi) {}_aN_b {}^be_{\beta}^{\text{s}\overline{\Gamma}}(\Phi),$$

and use these to obtain the **Field-Space Super-Metric**:

$${}_{\alpha}G_{\beta}(\Phi) = {}_{\alpha}e^a(\Phi) {}_aH_b {}^be_{\beta}^{\text{s}\overline{\Gamma}}(\Phi)$$

– The **Christoffel Symbols**

$${}^{\alpha}\Gamma_{\beta\gamma} = \frac{1}{2} G^{\alpha\delta} \left[{}_{\delta}G_{\beta} \overleftarrow{\partial}_{\gamma} + (-1)^{\beta\gamma} {}_{\delta}G_{\gamma} \overleftarrow{\partial}_{\beta} - (-1)^{\beta} \overrightarrow{\partial}_{\delta} {}_{\beta}G_{\gamma} \right]$$

– The **Riemann Tensor**

$$\begin{aligned} R^{\alpha}_{\beta\gamma\delta} = & - {}^{\alpha}\Gamma_{\beta\gamma} \overleftarrow{\partial}_{\delta} + (-1)^{\gamma\delta} {}^{\alpha}\Gamma_{\beta\delta} \overleftarrow{\partial}_{\gamma} + (-1)^{\gamma(\beta+\epsilon)} {}^{\alpha}\Gamma_{\epsilon\gamma} {}^{\epsilon}\Gamma_{\beta\delta} \\ & - (-1)^{\delta(\epsilon+\beta+\gamma)} {}^{\alpha}\Gamma_{\epsilon\delta} {}^{\epsilon}\Gamma_{\beta\gamma} \end{aligned}$$

- **Supergeometric Effective Action with Fermions**

[K Finn, S Karamitsos, AP, EPJC81 (2021) 572]

$$\exp(i\Gamma[\Phi]) = \int [D\Phi_q] \sqrt{|\text{sdet}G|} \exp \left(iS[\Phi_q] + i \int d^4x \sqrt{-g} \Gamma[\Phi] \frac{\overleftarrow{\partial}}{\partial \Phi^\alpha} \Sigma^\alpha[\Phi, \Phi_q] \right)$$

• Supergeometric Effective Action with Fermions

[K Finn, S Karamitsos, AP, EPJC81 (2021) 572]

$$\exp(i\Gamma[\Phi]) = \int [D\Phi_q] \sqrt{|\text{sdet} G|} \exp \left(iS[\Phi_q] + i \int d^4x \sqrt{-g} \Gamma[\Phi] \frac{\overleftarrow{\partial}}{\partial \Phi^\alpha} \Sigma^\alpha[\Phi, \Phi_q] \right)$$

– One- and Two-Loop Grand Covariant Effective Actions

$$\Gamma^{(1)}[\Phi] = \frac{i}{2} \ln \text{sdet} \left(\overrightarrow{\nabla}^{\hat{\alpha}} S \overleftarrow{\nabla}_{\hat{\beta}} \right) = \frac{i}{2} \text{str} \ln \left(\overrightarrow{\nabla}^{\hat{\alpha}} S \overleftarrow{\nabla}_{\hat{\beta}} \right)$$

$$\begin{aligned} \Gamma^{(2)}[\Phi] &= \text{Diagram 1} + \text{Diagram 2} \\ &= -\frac{1}{8} S \overleftarrow{\nabla}_{\hat{\alpha}} \overleftarrow{\nabla}_{\hat{\beta}} \overleftarrow{\nabla}_{\hat{\gamma}} \overleftarrow{\nabla}_{\hat{\delta}} \Delta^{\hat{\delta}\hat{\gamma}} \Delta^{\hat{\beta}\hat{\alpha}} \\ &\quad + (-1)^{\hat{\gamma}\hat{\beta} + \hat{\epsilon}(\hat{\delta} + \hat{\beta})} \frac{1}{12} \left(S \overleftarrow{\nabla}_{\hat{\epsilon}} \overleftarrow{\nabla}_{\hat{\gamma}} \overleftarrow{\nabla}_{\hat{\alpha}} \right) \Delta^{\hat{\alpha}\hat{\beta}} \Delta^{\hat{\gamma}\hat{\delta}} \Delta^{\hat{\epsilon}\hat{\zeta}} \left(\overrightarrow{\nabla}_{\hat{\zeta}} \overrightarrow{\nabla}_{\hat{\delta}} \overrightarrow{\nabla}_{\hat{\beta}} S \right) \end{aligned}$$

$$\hat{\alpha} \Delta_{\hat{\beta}}^{-1} \equiv \overrightarrow{\nabla}_{\hat{\alpha}} S \overleftarrow{\nabla}_{\hat{\beta}}, \text{ with } \hat{\alpha} \Delta^{\hat{\beta}} = \Delta^{\hat{\alpha}\hat{\beta}} \text{ defined through: } \hat{\alpha} \Delta^{\hat{\gamma}}_{\hat{\gamma}} \Delta_{\hat{\beta}}^{-1} = \hat{\alpha} \delta_{\hat{\beta}}.$$

– Single Fermion Model

$$\begin{aligned}\mathcal{L} = & \frac{1}{2} k(\phi) \partial_\mu \phi \partial^\mu \phi + \frac{i}{2} g(\phi) \left(\bar{\psi} \gamma^\mu \partial_\mu \psi - \partial_\mu \bar{\psi} \gamma^\mu \psi \right) \\ & - \frac{1}{2} h(\phi) \bar{\psi} \gamma^\mu \psi \partial_\mu \phi - Y(\phi) \bar{\psi} \psi - V(\phi)\end{aligned}$$

Supermanifold: $\Phi^\alpha = (\phi, \psi, \bar{\psi})$, with **grand** field-space metric

$${}_\alpha G_\beta = \begin{pmatrix} k - \frac{g'^2 + h^2}{2g} \bar{\psi} \psi & -\frac{1}{2}(g' - ih) \bar{\psi} & \frac{1}{2}(g' + ih) \psi \\ \frac{1}{2}(g' - ih) \bar{\psi} & 0_4 & g 1_4 \\ -\frac{1}{2}(g' + ih) \psi & -g 1_4 & 0_4 \end{pmatrix}$$

But, $R^\alpha_{\beta\gamma\delta} = 0 \implies$ field-space is **flat**

– **Frame**-reparametrization to a **Cartesian Frame**, $\tilde{\Phi}^\alpha = (\tilde{\phi}, \tilde{\psi}, \tilde{\bar{\psi}})^\top$:

$$\phi \rightarrow \tilde{\phi} = \int_0^\phi \sqrt{k(\phi)} d\phi, \quad \psi \rightarrow \tilde{\psi} = \sqrt{g(\phi)} \exp\left(\frac{i}{2} \int_0^\phi \frac{h(\phi)}{g(\phi)} d\phi\right) \psi$$

Lagrangian in the **Cartesian Frame**:

$$\mathcal{L} = \frac{1}{2} \partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi} + \frac{i}{2} \left(\tilde{\bar{\psi}} \gamma^\mu \partial_\mu \tilde{\psi} - \partial_\mu \tilde{\bar{\psi}} \gamma^\mu \tilde{\psi} \right) - \tilde{Y}(\tilde{\phi}) \tilde{\bar{\psi}} \tilde{\psi} - \tilde{V}(\tilde{\phi}),$$

with $\tilde{Y}(\tilde{\phi}) = Y(\phi)/g(\phi)$ and $\tilde{V}(\tilde{\phi}) = V(\phi)$.

– **Frame-reparametrization to a Cartesian Frame**, $\tilde{\Phi}^\alpha = (\tilde{\phi}, \tilde{\psi}, \tilde{\bar{\psi}})^\top$:

$$\phi \rightarrow \tilde{\phi} = \int_0^\phi \sqrt{k(\phi)} d\phi, \quad \psi \rightarrow \tilde{\psi} = \sqrt{g(\phi)} \exp\left(\frac{i}{2} \int_0^\phi \frac{h(\phi)}{g(\phi)} d\phi\right) \psi$$

Lagrangian in the **Cartesian Frame**:

$$\mathcal{L} = \frac{1}{2} \partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi} + \frac{i}{2} \left(\tilde{\bar{\psi}} \gamma^\mu \partial_\mu \tilde{\psi} - \partial_\mu \tilde{\bar{\psi}} \gamma^\mu \tilde{\psi} \right) - \tilde{Y}(\tilde{\phi}) \tilde{\bar{\psi}} \tilde{\psi} - \tilde{V}(\tilde{\phi}),$$

with $\tilde{Y}(\tilde{\phi}) = Y(\phi)/g(\phi)$ and $\tilde{V}(\tilde{\phi}) = V(\phi)$.

– **One-Loop Effective Action**

$$\begin{aligned} \Gamma[\Phi] = & S[\Phi] + \frac{i}{2} \text{Tr} \ln \left\{ \square + \tilde{V}'' - \tilde{\bar{\psi}} \left[2 \tilde{Y}'^2 (-i\not{\partial} + \tilde{Y})^{-1} - \tilde{Y}'' \right] \tilde{\psi} \right\} \\ & - i \text{Tr} \ln (-i\not{\partial} + \tilde{Y}) \end{aligned}$$

– **Model with Multiple Fermions** (up to quadratic kinetic terms)

$$\begin{aligned} \mathcal{L} = & \frac{1}{2} g^{\mu\nu} k_{AB}(\Phi) \partial_\mu \phi^A \partial_\nu \phi^B + \frac{i}{2} g_{XY}(\Phi) \left(\bar{\psi}^X \gamma^\mu \partial_\mu \psi^Y - \partial_\mu \bar{\psi}^X \gamma^\mu \psi^Y \right) \\ & - \frac{1}{2} h_{AXY}(\Phi) \bar{\psi}^X \gamma^\mu \psi^Y \partial_\mu \phi^A + \dots - U(\Phi) \end{aligned}$$

– **No-Go Theorem for Fermionic Curvature** [V Gattus, AP, arXiv:2307.01126]

Field-space super-metric for single scalar ϕ :

$${}_\alpha G_\beta = \begin{pmatrix} k - \frac{1}{2} \bar{\psi} (g' - i\mathbf{h}) g^{-1} (g' + i\mathbf{h}) \psi & -\frac{1}{2} \bar{\psi} (g' - i\mathbf{h}) & \frac{1}{2} \psi^\top (g'^\top + i\mathbf{h}^\top) \\ \frac{1}{2} (g'^\top - i\mathbf{h}^\top) \bar{\psi}^\top & \mathbf{0} & g^\top \mathbf{1}_4 \\ -\frac{1}{2} (g' + i\mathbf{h}) \psi & -g \mathbf{1}_4 & \mathbf{0} \end{pmatrix},$$

with $\psi = \{\psi^X\}$, $g(\phi) = \{g_{XY}\} = g^\dagger(\phi)$ and $\mathbf{h}(\phi) = \{h_{XY}\} = \mathbf{h}^\dagger(\phi)$.

But, $R^\alpha_{\beta\gamma\delta} = 0 \implies$ field super-space has zero curvature.

– Minimal 2D Model with Fermionic Curvature

[V Gattus, AP, arXiv:2307.01126]

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi + \frac{i}{2} \left(g_0(\phi) + \underline{g_1(\phi) \bar{\psi} \psi} \right) \left(\bar{\psi} \gamma^\mu \partial_\mu \psi - \partial_\mu \bar{\psi} \gamma^\mu \psi \right) - Y(\phi) \bar{\psi} \psi - V(\phi), \quad \text{with } \gamma^\mu = (\sigma^1, -i\sigma^2).$$

Chart: $\Phi = (\phi, \psi^\top, \bar{\psi})$, $\psi^\top = (\psi_1, \psi_2)$ and $\bar{\psi} \equiv \psi^\dagger \gamma^0$.

Field-space super-metric:

$${}_\alpha G_\beta = \begin{pmatrix} 1 + \mathbf{b}^\top (\mathbf{d}^{-1})^\top \mathbf{a}^\top - \mathbf{a} \mathbf{d}^{-1} \mathbf{b} & -\mathbf{a} & \mathbf{b}^\top \\ \mathbf{a}^\top & 0 & \mathbf{d}^\top \\ -\mathbf{b} & -\mathbf{d} & 0 \end{pmatrix},$$

where $\mathbf{d} = (g_0 + g_1 \bar{\psi} \psi) \mathbf{1}_2 + g_1 \psi \bar{\psi}$,

$\mathbf{a} = \bar{\psi} (g'_0 + g'_1 \bar{\psi} \psi)$, $\mathbf{b} = (g'_0 + g'_1 \bar{\psi} \psi) \psi$ (with $g'_{0,1} \equiv \partial g_{0,1} / \partial \phi$),

Set $g_0(\phi) = 1$ for simplicity:

Super-Ricci scalar: $\mathfrak{R} = 4g_1 - \frac{1}{2} g_1'^2 (\bar{\psi} \psi)^2 \neq 0 \implies$ field super-space is curved.

• Conclusions

- Complete formulation of **Geometric Effective Actions** for **Scalar–Tensor Theories**, with a **complete set** of model functions
- **New** model function $\ell = \ell(\Phi)$ ensures the **uniqueness** of the **VDW** path-integral measure, **with** $ds^2 = g_{\mu\nu}/\ell^2 dx^\mu dx^\nu$
- **Extension** of the **VDW** formalism to **Supermanifolds** to describe **realistic** theories that include **fermions**, such as the SM
- **Minimal Supergeometric QFT Models** with **non-zero fermionic curvature** [See talk by V Gattus]
- **Analytic expressions** of **One- and Two-Loop Supergeometric Effective Actions** for **Theories with Fermions**

• Future Research Directions

- UV completions of Minimal Supergeometric QFTs (Plan A)
(avoiding the Haag-Lopuszanski-Sohnius theorem)
- Application of Supergeometric QFTs to SMEFT and HEFT (Plan B)
- Include gauge and other symmetries $\iff$ Isometries on Supermanifold
- Phenomenological/cosmological applications of Supergeometric QFTs

⋮

Supergeometry on a Supermanifold

$$D\Sigma^2 = d\Phi^{\hat{\alpha}} \, {}_{\hat{\alpha}}G_{\hat{\beta}}(\Phi) \, d\Phi^{\hat{\beta}}$$


[K Finn, S Karamitsos, AP, [EPJC81 \(2021\) 572](#);
Viola Gattus, AP, [arXiv:2307.01126](#)]

Supergeometry on a Supermanifold

$$D\Sigma^2 = d\Phi^{\hat{\alpha}} \hat{\alpha} G_{\hat{\beta}}(\Phi) d\Phi^{\hat{\beta}}$$

[K Finn, S Karamitsos, AP, [EPJC81 \(2021\) 572](#);
Viola Gattus, AP, [arXiv:2307.01126](#).]

from Plato



Back-Up Slides

- From **Geometrizing** the **Cosmos** to **Micro-Cosmos**

- **Geometrizing the Cosmos**: . . . , Pythagoras (5c BC)

- **Geocentric versus Heliocentric System** [e.g., Van der Waerden '87]

Geocentric: Anaximander (6c BC), . . . , Plato (4c BC),
Aristotle (3c BC), Ptolemy (2c AD), . . . **Tycho (16c AD)**

Heliocentric: Aristarchus (3c BC), Seleucus (2c BC),
Copernicus (15c AD), Kepler (16c AD), Galileo (16c AD), . . .

- **Absolute versus Relative/Local Inertial Frame in Gravitation**

Absolute: Newton (17c AD), . . .

Relative: Einstein (20c AD), . . .


- **Geometrizing the Micro-Cosmos** as a solution to frame problems in Quantum Field Theory and Quantum Gravity

– Einstein versus Jordan Frame

Action in **Einstein Frame**: $S^{\text{EF}}[g_{\mu\nu}, \varphi] = \int_x \left[-\frac{1}{2}M_P^2 R + \frac{1}{2}(\partial_\mu \varphi)^2 - V(\varphi) \right]$

Action in **Jordan Frame**: $S^{\text{JF}}[\tilde{g}_{\mu\nu}, \tilde{\varphi}] = \int_x \left[-\frac{1}{2}f(\tilde{\varphi})\tilde{R} + \frac{1}{2}(\partial_\mu \tilde{\varphi})^2 - \tilde{V}(\tilde{\varphi}) \right]$

Frame equivalence $\implies S^{\text{JF}}[\tilde{g}_{\mu\nu}, \tilde{\varphi}] = S^{\text{EF}}[g_{\mu\nu}, \varphi]$ [R. H. Dicke '62]



$\Gamma^{\text{JF}}_{1\text{-loop}}[\tilde{g}_{\mu\nu}, \tilde{\varphi}] \neq \Gamma^{\text{EF}}_{1\text{-loop}}[g_{\mu\nu}, \varphi]$: **Effective action is frame dependent, except at extrema of the action.**

– Partial list of references

- R. H. Dicke, *Mach's principle and invariance under transformation of units*, Phys. Rev. 125 (1962) 2163.
- V. Faraoni, E. Gunzig, P. Nardone, *Conformal transformations in classical gravitational theories and in cosmology*, Fund. Cosmic Phys. 20 (1999) 121.
- E. E. Flanagan, *The Conformal Frame Freedom in Theories of Gravitation*, Class. Quant. Grav. 21 (2004) 3817.
- T. Chiba, M. Yamaguchi, *Conformal-Frame (In)dependence of Cosmological Observations in Scalar-Tensor Theory*, JCAP 1310 (2013) 040.
- M. Postma, M. Volponi, *Equivalence of the Einstein and Jordan frames*, Phys. Rev. D90 (2014) 103516.
- A. Y. Kamenshchik, C. F. Steinwachs, *Question of quantum equivalence between Jordan frame and Einstein frame*, Phys. Rev. D91 (2015) 084033.
- D. Burns, S. Karamitsos, A. Pilaftsis, *Frame-Covariant Formulation of Inflation in Scalar-Curvature Theories*, Nucl. Phys. B907 (2016) 785.
- L. Järv, K. Kannike, L. Marzola, A. Racioppi, M. Raidal, M. Rnkla, M. Saal, H. Veerme, *Frame-Independent Classification of Single-Field Inflationary Models*, Phys. Rev. Lett. 118 (2017) 151302.
- G. Domnech, M. Sasaki, *Conformal Frame Dependence of Inflation*, JCAP 1504 (2015) 022.
- J. O. Gong, T. Tanaka, *A covariant approach to general field space metric in multi-field inflation*, JCAP 1103 (2011) 015, Erratum: [JCAP 1202 (2012) E01].
- A. Karam, T. Pappas, K. Tamvakis, *Frame-dependence of higher-order inflationary observables in scalar-tensor theories*, Phys. Rev. D96 (2017) 064036.
- S. Karamitsos and A. Pilaftsis, *Frame Covariant Nonminimal Multifield Inflation*, Nucl. Phys. B927 (2018) 219.

- Effects of $\ell(\varphi)$ (from $\bar{g}_{\mu\nu} \equiv g_{\mu\nu}/\ell^2(\varphi)$)

- Frame-invariant **Dirac-delta** function

$$\int d^D x \sqrt{-\bar{g}} \bar{\delta}^{(D)}(x) = 1, \quad \text{with} \quad \bar{\delta}^{(D)}(x_I - x_J) \equiv \frac{\delta^{(D)}(x_I - x_J)}{\sqrt{-\bar{g}}}$$

- Functional **derivative**

$$\frac{\bar{\delta} F[\Phi(x)]}{\bar{\delta} \Phi(y)} \equiv \lim_{\epsilon \rightarrow 0} \frac{F[\Phi(x) + \epsilon \bar{\delta}^{(D)}(x - y)] - F[\Phi(x)]}{\epsilon}$$

- Functional **determinant**

$$\overline{\det}(M_{xy}) \equiv \exp \left[i \int d^D x \sqrt{-\bar{g}} \ln(M)_{xx} \right]$$

• **Field-Space Riemann Tensor** $\mathfrak{R}^{(\mu\nu)}_{(\alpha\beta)(\rho\sigma)(\gamma\delta)}$ **for General Relativity**

[K Finn, S Karamitsos, AP, '19]

$$\begin{aligned}
 \mathfrak{R}^{(\mu\nu)}_{(\alpha\beta)(\rho\sigma)(\gamma\delta)} = & -\frac{1}{32}\delta_\rho^\mu\delta_\beta^\nu g_{\sigma\gamma}g_{\alpha\delta} - \frac{1}{32}\delta_\sigma^\mu\delta_\beta^\nu g_{\rho\gamma}g_{\alpha\delta} - \frac{1}{32}\delta_\beta^\mu\delta_\sigma^\nu g_{\rho\gamma}g_{\alpha\delta} - \frac{1}{32}\delta_\beta^\mu\delta_\rho^\nu g_{\sigma\gamma}g_{\alpha\delta} \\
 & -\frac{1}{32}\delta_\rho^\mu\delta_\beta^\nu g_{\sigma\delta}g_{\alpha\gamma} - \frac{1}{32}\delta_\sigma^\mu\delta_\beta^\nu g_{\rho\delta}g_{\alpha\gamma} - \frac{1}{32}\delta_\beta^\mu\delta_\sigma^\nu g_{\rho\delta}g_{\alpha\gamma} - \frac{1}{32}\delta_\beta^\mu\delta_\rho^\nu g_{\sigma\delta}g_{\alpha\gamma} \\
 & -\frac{1}{32}\delta_\alpha^\mu\delta_\rho^\nu g_{\sigma\gamma}g_{\beta\delta} - \frac{1}{32}\delta_\alpha^\mu\delta_\sigma^\nu g_{\rho\gamma}g_{\beta\delta} - \frac{1}{32}\delta_\rho^\mu\delta_\alpha^\nu g_{\sigma\gamma}g_{\beta\delta} - \frac{1}{32}\delta_\sigma^\mu\delta_\alpha^\nu g_{\rho\gamma}g_{\beta\delta} \\
 & -\frac{1}{32}\delta_\alpha^\mu\delta_\rho^\nu g_{\sigma\delta}g_{\beta\gamma} - \frac{1}{32}\delta_\alpha^\mu\delta_\sigma^\nu g_{\rho\delta}g_{\beta\gamma} - \frac{1}{32}\delta_\rho^\mu\delta_\alpha^\nu g_{\sigma\delta}g_{\beta\gamma} - \frac{1}{32}\delta_\sigma^\mu\delta_\alpha^\nu g_{\rho\delta}g_{\beta\gamma} \\
 & +\frac{1}{32}\delta_\gamma^\mu\delta_\beta^\nu g_{\rho\delta}g_{\sigma\alpha} + \frac{1}{32}\delta_\delta^\mu\delta_\beta^\nu g_{\rho\gamma}g_{\sigma\alpha} + \frac{1}{32}\delta_\beta^\mu\delta_\delta^\nu g_{\rho\gamma}g_{\sigma\alpha} + \frac{1}{32}\delta_\beta^\mu\delta_\gamma^\nu g_{\rho\delta}g_{\sigma\alpha} \\
 & +\frac{1}{32}\delta_\gamma^\mu\delta_\beta^\nu g_{\rho\alpha}g_{\sigma\delta} + \frac{1}{32}\delta_\delta^\mu\delta_\beta^\nu g_{\rho\alpha}g_{\sigma\gamma} + \frac{1}{32}\delta_\beta^\mu\delta_\delta^\nu g_{\rho\alpha}g_{\sigma\gamma} + \frac{1}{32}\delta_\beta^\mu\delta_\gamma^\nu g_{\rho\alpha}g_{\sigma\delta} \\
 & +\frac{1}{32}\delta_\alpha^\mu\delta_\gamma^\nu g_{\rho\delta}g_{\sigma\beta} + \frac{1}{32}\delta_\alpha^\mu\delta_\delta^\nu g_{\rho\gamma}g_{\sigma\beta} + \frac{1}{32}\delta_\gamma^\mu\delta_\alpha^\nu g_{\rho\delta}g_{\sigma\beta} + \frac{1}{32}\delta_\delta^\mu\delta_\alpha^\nu g_{\rho\gamma}g_{\sigma\beta} \\
 & +\frac{1}{32}\delta_\alpha^\mu\delta_\gamma^\nu g_{\rho\beta}g_{\sigma\delta} + \frac{1}{32}\delta_\alpha^\mu\delta_\delta^\nu g_{\rho\beta}g_{\sigma\gamma} + \frac{1}{32}\delta_\gamma^\mu\delta_\alpha^\nu g_{\rho\beta}g_{\sigma\delta} + \frac{1}{32}\delta_\delta^\mu\delta_\alpha^\nu g_{\rho\beta}g_{\sigma\gamma} \\
 & +\frac{1}{4D}g_{\rho\gamma}g^{\mu\nu}g_{\sigma\beta}g_{\alpha\delta} + \frac{1}{4D}g_{\rho\delta}g^{\mu\nu}g_{\sigma\beta}g_{\alpha\gamma} + \frac{1}{4D}g_{\rho\alpha}g^{\mu\nu}g_{\sigma\gamma}g_{\beta\delta} + \frac{1}{4D}g_{\rho\alpha}g^{\mu\nu}g_{\sigma\delta}g_{\beta\gamma} \\
 & +\frac{1}{4D}g_{\rho\gamma}g^{\mu\nu}g_{\sigma\alpha}g_{\beta\delta} + \frac{1}{4D}g_{\rho\delta}g^{\mu\nu}g_{\sigma\alpha}g_{\beta\gamma} + \frac{1}{4D}g_{\rho\beta}g^{\mu\nu}g_{\sigma\delta}g_{\alpha\gamma} + \frac{1}{4D}g_{\rho\beta}g^{\mu\nu}g_{\sigma\gamma}g_{\alpha\delta} \\
 & -\frac{1}{4D}g^{\mu\nu}g_{\rho\beta}g_{\sigma\gamma}g_{\alpha\delta} - \frac{1}{4D}g^{\mu\nu}g_{\rho\beta}g_{\sigma\delta}g_{\alpha\gamma} - \frac{1}{4D}g^{\mu\nu}g_{\rho\gamma}g_{\sigma\alpha}g_{\beta\delta} - \frac{1}{4D}g^{\mu\nu}g_{\rho\delta}g_{\sigma\alpha}g_{\beta\gamma} \\
 & -\frac{1}{4D}g^{\mu\nu}g_{\rho\alpha}g_{\sigma\gamma}g_{\beta\delta} - \frac{1}{4D}g^{\mu\nu}g_{\rho\alpha}g_{\sigma\delta}g_{\beta\gamma} - \frac{1}{4D}g^{\mu\nu}g_{\rho\delta}g_{\sigma\beta}g_{\alpha\gamma} - \frac{1}{4D}g^{\mu\nu}g_{\rho\gamma}g_{\sigma\beta}g_{\alpha\delta}
 \end{aligned}$$

– Field-Space Ricci Tensor $\mathfrak{R}_{(\mu\nu)(\rho\sigma)}$

$$\mathfrak{R}_{(\mu\nu)(\rho\sigma)} = \frac{1}{4}g_{\mu\nu}g_{\rho\sigma} - \frac{D}{8}g_{\mu\rho}g_{\nu\sigma} - \frac{D}{8}g_{\mu\sigma}g_{\nu\rho}$$

– Field-Space Ricci Scalar $\mathfrak{R}$

$$\mathfrak{R} = \frac{D}{4} - \frac{D^2}{8} - \frac{D^3}{8} < 0,$$

for spacetime dimensions $D > 1$.

$\Rightarrow$ Gravity has a curved field space, with *negative* scalar curvature.

- **Partial differentiation with respect to γ^μ**

Any matrix $M_{a\dot{a}}$ in spinor space can be uniquely expressed as

$$M_{a\dot{a}} = \sum_{i=S,P,V,A,T} a_{(i)} \Gamma_{a\dot{a}}^{(i)}, \quad \text{with } \Gamma^{(i)} \in \{I_4, \gamma^5, \gamma^\mu, \gamma^\mu \gamma^5, \sigma^{\mu\nu}\},$$

where $\Gamma^{(i)}$ are the 16 orthogonal Lorentz-covariant bilinears, $a_{(i)}$ is a coefficient, and $\sigma^{\mu\nu} \equiv i/2 [\gamma^\mu, \gamma^\nu]$.

- **Γ -matrix partial differentiation:**

$$\frac{\delta M}{\delta \Gamma^{(i)}} \equiv \lim_{\epsilon^{(i)} \rightarrow 0} \frac{M[\Gamma^{(i)} \rightarrow \Gamma^{(i)} + \epsilon^{(i)} I_4] - M[\Gamma^{(i)}]}{\epsilon^{(i)}}.$$

For $\Gamma^{(i)} = \Gamma^{(V)} = \gamma^\mu$,

$$\frac{\delta M_{a\dot{a}}}{\delta \gamma^\mu} = a_\mu \delta_{a\dot{a}}.$$