

The Hunt for Extra Higgs Bosons in the NMSSM

Ulrich Ellwanger
Université Paris-Saclay, Orsay, France
with Matías Rodríguez-Vázquez



December 3, 2017

The NMSSM: MSSM + an extra Singlet (Super-)Field \widehat{S}

- Solves the μ -problem of the MSSM:
 $\langle S \rangle$ generates the required higgsino mass term
- Alleviates the tension related to the SM-like Higgs mass:
 - In the MSSM the tree level quartic Higgs coupling is bounded by the electroweak gauge couplings $\rightarrow M_H \leq M_Z \rightarrow$ need large radiative corrections
 - The extra singlet in the NMSSM generates an extra quartic Higgs coupling $\sim \lambda^2 \rightarrow$ less radiative corrections required
- The “singlino” is a perfect dark matter candidate, good relic density compatible with its non-observation in direct and indirect searches

The Higgs Sector of the NMSSM: ("Higgs basis")

- An essentially SM-like H_{125} , up to mixing with H_{MSSM}/H_S
- H_{MSSM} , a heavy approximately degenerate SU(2) doublet:
neutral scalar + pseudoscalar + one charged state;
heavy due to lower bounds on M_{H^\pm} amongst others from $b \rightarrow s + \gamma$
- Essentially singlet-like scalar H_S and pseudoscalar A_S ,
NOT degenerate in general, lighter or heavier than 125 GeV
depending on unknown parameters


How to Search for H_S/A_S : Direct Production:

Rely on mixing $\sim \xi$ of H_S with H_{125} : Induces couplings of H_S to SM particles
→ same production mechanisms but with smaller cross sections

Mixing reduces these couplings of H_{125} → limited to $\xi^2 \lesssim 0.25$ (at present)

Summary of LEP search for a light scalar
decaying into $b\bar{b}$
with reduced coupling ξ^2 to ZZ :



The region in the $\xi^2 - m_H$ plane
below the black line is allowed
→ Weak bounds for $M_{H_S} \sim 90 - 100$ GeV



ATLAS and CMS searches for $ggF \rightarrow H_S \rightarrow \gamma\gamma$ at 8 TeV:

ATLAS, PRL 113 (2014) 17, 171801, 1407.6583

CMS-PAS-HIG-14-037




CMS search for $ggF \rightarrow H_S \rightarrow \gamma\gamma$ at 13 TeV:

(CMS PAS HIG-17-013)

→ Excesses near 97 GeV


(not confirmed by ATLAS)



Do the 8 TeV ATLAS/CMS searches touch possible values for $\sigma(\text{gg}F \rightarrow H_S \rightarrow \gamma\gamma)$ within the LEP-allowed NMSSM parameter space?

(M. Rodríguez, U.E., JHEP 1602 (2016) 096):

Red points: NMSSM-specific lift of $\Delta M_{H_{125}} > 12$ GeV




YES, the excess could be explained (if confirmed) ... but even light H_S states may have too small direct production cross sections for discovery, even at 13 TeV

Run I searches for $H_{125} \rightarrow A_5 A_5 (H_5 H_5) \rightarrow 4$ leptons

(From R. Aggleton et al., JHEP 1702 (2017) 035, arXiv:1609.06089)

Observed exclusion limits ($\sqrt{s} = 8$ TeV)



Light green/blue points: viable in the NMSSM after LEP/LHC constraints

→ These searches for H_5/A_5 have only scratched the NMSSM parameter space ...

... and are limited to $M_{H_S, A_S} \lesssim 60$ GeV; how to search for heavier H_S/A_S ?

→ Note: the trilinear couplings $H_{MSSM}H_S H_{125}$, $A_{MSSM}A_S H_{125}$ are SU(2) invariant and can be large (proportional to a parameter A_λ), in contrast to $H_{MSSM}H_{125}H_{125}$, $H_{MSSM}H_S H_S$, ... which must be proportional to a SU(2) breaking vev \times couplings

→ The BRs($H_{MSSM}/A_{MSSM} \rightarrow H_{125}H_S/A_S$) can be large $\sim 50\%$, competing only with $H/A_{MSSM} \rightarrow t\bar{t}$, and reducing the BR for the search into $\tau\tau$!

$ggF \rightarrow H_{MSSM} \rightarrow H_S H_{125}$ and $ggF \rightarrow A_{MSSM} \rightarrow A_S H_{125}$ look like resonant Higgs pair production, but with one SM Higgs replaced by H_S/A_S with unknown mass

A New Search: Look for $b\bar{b}b\bar{b}$ or $b\bar{b}\gamma\gamma$

with

- one $b\bar{b}$ or $\gamma\gamma$ pair: $M_{b\bar{b}/\gamma\gamma} \sim 125$ GeV,
- another $b\bar{b}$ or $\gamma\gamma$ pair: $M_{b\bar{b}/\gamma\gamma} \sim M_{H_S, A_S}$ (unknown),
- $M_{b\bar{b}b\bar{b}}$ or $M_{b\bar{b}\gamma\gamma} \sim M_{H/A_{MSSM}}$ (unknown)

$b\bar{b}\tau\tau$ final states are slightly less promising;

Best Strategy (M. Rodríguez, U.E., JHEP_072P_0817)

Borrowed from ATLAS/CMS searches for resonant SM Higgs pair production:

Use a “test” mass M_{H_5} ; for given M_{H_5} (or M_{A_5}):

- Optimize the cuts on p_T ,
- Optimise the pairing of 4 b -tagged jets into 2×2 b -tagged jets: cut on bb masses around 115 GeV and $M_{H_5} - 10$ GeV (allow for “losses” outside the $R = 0.4$ – jets)
- Study the distribution of M_{4b} from the 4 b -tagged jets (after correcting both $M_{b\bar{b}}$ to 125 GeV/ M_{H_5})
- Only if M_{H_5} was chosen correctly, one observes a “bump” in $M_{b\bar{b}b\bar{b}}$ near $M_{H_{MSSM}}$ whose significance can be computed as function of M_{H_5} , $M_{H_{MSSM}}$ and notably the $\sigma(ggF \rightarrow H_{MSSM} \rightarrow H_{125} + H_5 \rightarrow b\bar{b}b\bar{b})$
- Expected 95% CL exclusion limits and 5σ discovery limits can be obtained
- These are model independent (assuming just a width \lesssim a few GeV)



Background





The dominant QCD backgrounds are multijets $b\bar{b}b\bar{b}$, $b\bar{b}c\bar{c}$ (mistagged), and a few % $t\bar{t}$. Simulations are insufficient; the absolute scale should be obtained from sidebands as done by ATLAS/CMS in their search for Higgs pair production in ATL-CONF-2016-049, CMS-PAS-HIG-16-002:

- Simulate multijets (+ $t\bar{t}$)
- Compare to data in sidebands (2 b-tags only)
- Rescale the simulated M_{4b} distribution into the signal region (4 b-tags)
- obtain a realistic background estimate

Here: Take $M_{H_S} \sim 125$ GeV, simulate multijets (+ $t\bar{t}$), apply ATLAS cuts, compare our simulated M_{4b} distribution to ATLAS data in their search for Higgs pair production, used as “sideband”

→ Rescale the simulated M_{4b} distribution from multijets by a factor 1.55 ± 0.27 (beyond a NLO K-factor 1.7): Approximately M_{4b} independent!



L.h.s.: M_{4b} distribution from MC vs. ATL-CONF-2016-049.

R.h.s.: M_{4b} distribution from MC vs. ATL-CONF-2016-049 after rescaling

Subsequently: Rescale M_{4b} distributions from MC by 1.55 ± 0.27 for all M_{H_S}

Exp. sensitiv. to $\sigma(ggF \rightarrow H/A \rightarrow H_{SM} + H_S/A_S \rightarrow b\bar{b}b\bar{b})$ as function of M_{H_S} :





Upper left: $M_{H_{MSSM}} = 500$ GeV, 95% limits and 5σ discovery for $L = 300fb^{-1}$

Upper right: $M_{H_{MSSM}} = 750$ GeV, 95% limits and 5σ discovery for $L = 300fb^{-1}$

Lower left: $M_{H_{MSSM}} = 500$ GeV, 95% limits and 5σ discovery for $L = 3000fb^{-1}$

Lower right: $M_{H_{MSSM}} = 750$ GeV, 95% limits and 5σ discovery for $L = 3000fb^{-1}$

Blue: NMSSM points for H_S , Orange: NMSSM points for A_S



Exp. sensitiv. to $\sigma(ggF \rightarrow H/A \rightarrow H_{SM} + H_S/A_S \rightarrow b\bar{b}\gamma\gamma)$ as function of M_{H_S} :





Upper left: $M_{H_{MSSM}} = 425$ GeV, 95% limits and 5σ discovery for $L = 300fb^{-1}$

Upper right: $M_{H_{MSSM}} = 500$ GeV, 95% limits and 5σ discovery for $L = 300fb^{-1}$

Lower left: $M_{H_{MSSM}} = 425$ GeV, 95% limits and 5σ discovery for $L = 3000fb^{-1}$

Lower right: $M_{H_{MSSM}} = 500$ GeV, 95% limits and 5σ discovery for $L = 3000fb^{-1}$

Blue: NMSSM points for H_S , Orange: NMSSM points for A_S



Conclusions

- The NMSSM contains a rich BSM Higgs sector which is hardly tested
- Mostly singlet-like scalars H_S (or A_S) are difficult to detect:
 - If lighter than 60 GeV possibly via H_{125} decays; indirect constraints from SM-like signal rates of H_{125} notably $H_{125} \rightarrow Z^*Z$, many ongoing ATLAS/CMS studies on exotic Higgs decays
 - If heavier than 60 GeV: $ggF \rightarrow H_S/A_S \rightarrow \gamma\gamma$ cross sections may be large enough if H_S/A_S mix considerably with H_{125} , H_{MSSM}/A_{MSSM} ; excesses near $M_{H_S} \sim 97$ GeV remain to be confirmed
- Searches for $H_{MSSM}/A_{MSSM} \rightarrow H_{125} + H_S/A_S$ can even be sensitive to pure singlet-like H_S/A_S !
(Such decays would also reduce the branching ratio of H_{MSSM}/A_{MSSM} into the usually employed $\tau\tau$ channel \rightarrow alleviated limits!)
Allow to discover simultaneously H_{MSSM}/A_{MSSM} and H_S/A_S !